



Tampa Bay ENT

Adult and Pediatric Ear, Nose and Throat • Facial Plastic Surgery

NASAL SINUS SURGERY
HEAD AND NECK SURGERY
FACIAL PLASTIC SURGERY
PEDIATRIC OTOLARYNGOLOGY
HEARING EVALUATIONS, HEARING AIDS
DIZZINESS EVALUATIONS

DENNIS S. AGLIANO, M.D., F.A.C.S.
RENE A. BOOTHBY, M.D.
MIGUEL A. RIVERA, M.D.
KARIN S. HOTCHKISS, M.D.
SCOTT A. POWELL, M.D., M.B.A.
KRISTEN H. DECELLES, Au.D.

SINUS SURGERY INSTRUCTIONS

1. Take Tylenol or the prescription you have been given for pain. If possible start *Afrin Nasal Spray* the evening of surgery and continue 1 spray each nostril twice a day for 3 days: then stop. This will help reduce the drainage after surgery. Also start *Nasal Saline/Salt Water* spray 3 times daily until your follow up visit.
2. **DO NOT** take aspirin or Ibuprofen products.
3. A certain amount of blood and/or blood tinged mucous is expected. Change the nasal drip pad as necessary. The drainage will improve over the first 3 days.
4. A small degree of swelling of the external nose or the upper lip may occur. Applying a light ice pack (crushed ice in a baggie) to reduce the swelling.
5. Keeping the head elevated at 45 degrees day and night for the first week will help with swelling.
6. Apply *Vaseline* or *Chap Stick* to the lips if they get dry.
7. Soft or liquid diet if there is any nausea after anesthesia.
8. Gently clean the nostrils with a Q-tip and hydrogen peroxide twice a day as needed to remove crusting.
9. **Do not blow or wipe your nose.** Sneeze with mouth open. You may dab at the nostrils gently with a tissue.
10. You should not swim for at least 2 weeks and no scuba diving for at least two months.
11. Be sure to rest as much as possible for the first 5-7 days.
12. Avoid any bending and lifting anything heavier than a gallon of milk for the first week.
13. Report to your doctor if:
 - a. Excessive pain and/or swelling
 - b. Temperature over 101° F
 - c. Any injury to your nose
 - d. Excessive bleeding